

Antibodies

A LABORATORY MANUAL

Ed Harlow

Cold Spring Harbor Laboratory

David Lane

Imperial Cancer Research Fund Laboratories



Cold Spring Harbor Laboratory
1988

R E C E I V E D
OCT 25 1989

TOWNSEND AND TOWNSEND P.A.
LIBRARY

Antibodies

A LABORATORY MANUAL

All rights reserved

© 1988 by Cold Spring Harbor Laboratory

Printed in the United States of America

Book and cover design by Emily Harste

Cover: "Nature Abstracted," watercolor by Carl Molno

Library of Congress Cataloging-in-Publication Data

Antibodies : a laboratory manual / by Ed Harlow, David Lane.

p. cm.
Bibliography: p.
Includes index.

ISBN 0-87969-314-2

1. Immunoglobulins--Laboratory manuals. 2. Immunochemistry--

Laboratory manuals. I. Harlow, Ed. II. Lane, David (David P.).

1952--

QR186.7.A53 1988

574.2'93'028--dc19

Researchers using the procedures of this manual do so at their own risk. Cold Spring Harbor Laboratory makes no representations or warranties with respect to the material set forth in this manual and has no liability in connection with the use of these materials.

Certain experimental procedures in this manual may be the subject of national or local legislation or agency restrictions. Users of this manual are responsible for obtaining the relevant permissions, certificates, or licenses in these cases. Neither the authors of this manual nor Cold Spring Harbor Laboratory assume any responsibility for failure of a user to do so.

Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted by Cold Spring Harbor Laboratory for libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that the base fee of \$1.00 per article is paid directly to CCC, 27 Congress St., Salem MA 05970. [0-87969-314-2/88 \$1.00 + .00]. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale.

All Cold Spring Harbor Laboratory publications may be ordered directly from Cold Spring Harbor Laboratory, Box 100, Cold Spring Harbor, New York 11724. Phone 1-800-843-4388. In New York (516) 367-8423.

The interaction of an antibody with an antigen forms the basis of all immunochemical techniques. This chapter discusses the properties of the antibody-antigen interaction and is divided into three sections. The first summarizes the structure of the antibody-antigen bonds, the second covers the strength of these interactions, a characteristic known as affinity, and the third presents the factors that contribute to the overall stability of immune complexes, a property called avidity.

STRUCTURE OF THE ANTIBODY-ANTIGEN COMPLEX

The structure of the antibody-antigen complex has been studied by measuring the affinity of binding between an antibody and a series of related antigens, by using affinity labeling reagents, by site-directed mutagenesis of the antibody combining site, by molecular modeling, and, most compellingly, by X-ray diffraction studies of antibody-antigen cocrystals. Together, these techniques have delineated the region of the antibody molecule that is involved in antigen binding, the region of the antigen molecule that interacts with the antibody, and the molecular basis for antibody specificity.

The antigen binding site of an antibody is formed by the variable regions of the heavy and light chains

Affinity labeling and X-ray crystallography of immune complexes have established that the antigen binding site is formed by the heavy- and light-chain variable regions (see Fig. 2.5). The two variable regions are closely associated and are bound to each other by noncovalent interactions. The remainder of the heavy and light chains forms other domains that are not involved in antigen binding (see Chapter 2). The amino acids forming the antigen binding site are derived from both the heavy and light chains and correspond to the amino acids of the hypervariable regions determined from protein sequencing. The hypervariable regions are known as the complementarity determining regions (CDRs). There are six CDRs, three on each chain, and they form discrete loops anchored and oriented by the framework residues of the variable domains (Fig. 3.1).

The region of an antigen that binds to an antibody is called an epitope

The region of an antigen that interacts with an antibody is defined as an epitope. An epitope is not an intrinsic property of any particular